



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C12N 15/57, 9/64, A61K 31/70, 38/48, C12Q 1/37, A61K 48/00, C07K 16/40, C12N 15/00, A01K 67/027, C12Q 1/68	A1	(11) International Publication Number: WO 98/49322 (43) International Publication Date: 5 November 1998 (05.11.98)
(21) International Application Number: PCT/IB98/00625 (22) International Filing Date: 24 April 1998 (24.04.98) (30) Priority Data: 966/97 26 April 1997 (26.04.97) CH (71)(72) Applicant and Inventor: SONDEREGGER, Peter [CH/CH]; Biochemisches Institut Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich (CH). (74) Agent: ZINK-WILD, Markus, Peter; Patentanwaltsbüro Zink, Birchlistrasse 11, CH-8173 Riedt-Neerach (CH).		(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, GW, HU, ID, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
(54) Title: NEUOTRYPSIN (57) Abstract There are described neurotrypsins of the formulas (I) or (II), including the separate coding and coded sequences of these compounds of the formulas (I) or (II). These compounds may be used as at least one active compound in a pharmaceutical component. The coded peptide sequences of these compounds may be used as targets for the development of pharmaceutical drugs.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Neurotrypsin

Technical Field

5

The present invention is directed to neurotrypsins and to a pharmaceutical composition which contains these substances or has an influence on these substances.

10 Disclosure of Invention

Neurotrypsin is a newly discovered serine protease, which is predominantly expressed in the brain and in the lungs; the expression in the brain takes place nearly exclusively in the neurons.

15

Neurotrypsin has a previously not yet found domain composition: besides the protease domain, there are found 3 or 4 SRCR (scavenger receptor cysteine-rich) domains and one Kringle domain. It is to be pointed out that the combination of Kringle and SRCR domains have not yet been found in proteins. At the amino terminus of the neurotrypsin protein there is a segment of more than 60 amino acids, which has an extremely high proportion of proline and basic amino acids (arginine and histidine).

20

The invention is characterized by the characteristics in the independent claims. Preferred embodiments are defined in the dependent claims.

25

The newly found neurotrypsins

- neurotrypsin of the human (compound of the formula I),
- neurotrypsin of the mouse (compound of the formula II)

30 differ structurally very much from the so far known serine proteases.

The serine protease whose protease domain is structurally most closely related with the protease domain of the new compounds, namely plasmin (of the human), has only a 44 % amino acid sequence identity.

35

Th proline-rich, basic segment at the amino terminus has a certain resemblance with th basic segments of the netrins and the semaphorins/collapsins. Due to this

segment, it is probable that neurotrypsin may be enriched by means of heparin-affinity chromatography.

5 The neurotrypsins of the human (compound of the formula I) and of the mouse (compound of the formula II) exhibit a very high structural similarity among each other.

The identity of the amino acid sequences of the native proteins of the compounds of the formulas I or II amounts to 81%.

10 The neurotrypsin of the human (compound of the formula I) has a coding sequence of 2625 nucleotides. The coded peptide of the compound of the formula I has a length of 875 amino acids and contains a signal peptide of 20 amino acids. The neurotrypsin of the mouse (compound of the formula II) has a coding sequence of 2283 nucleotides. The coded protein of the compound of the formula II has a length of 761
15 amino acids and contains a signal peptide of 21 amino acids. The reason for the greater length of the neurotrypsin of the human consists therein that the human neurotrypsin has 4 SRCR domains, whereas the neurotrypsin of the mouse has only 3 SRCR domains.

20 The domains which are present in both compounds (compound of the formula I and compound of the formula II) have a high degree of sequence similarity. The corresponding SRCR domains of the compounds of the formulas I and II have an amino acid sequence identity from 81% to 91%. The corresponding Kringle domains have an amino acid sequence identity of 75%. A high degree of similarity consists also in the enzymatically active (i.e. proteolytic) domain (90% amino acid sequence identity).

25

The protease domains of the neurotrypsins of the human (compound of the formula I) and of the mouse (compound of the formula II) are aligned in the following section, in order to illustrate the high degree of sequence identity.

CGLRLLHRRQKRIIGGKNSLRGGWPWQVSLRLKSSHGDGRLLCGATLLSS : . : . CGLRLLHRRQKRIIGGNSLRGAWPWQASLRLRSAHGDGRLLCGATLLSS	50
CWVLTAAHCFKRYGNSTRSYAVRVGDYHTLVPEEFEEEEIGVQQIVIHREY : . : : CWVLTAAHCFKRYGNNSRSYAVRVGDYHTLVPEEFEEQIGVQQIVIHREY	100
RPDRSDYDIALVRLQGPEEQCARFSSHVLPACLPWRERPQKTASNCYIT : : . : . RPDRSDYDIALVRLQGPGEQCARLSTHVLPACLPLWRERPQKTASNCHIT	150
GWGDTGRAYSR TLQQAAPLLPKRFCEERYKGRFTGRMLCAGNLHEHKRV : . : . GWGDTGRAYSR TLQQAAPLLPKRFCKERYKGLFTGRMLCAGNLQEDNRV	200
DSCQGDSSGGPLMCERPGE SWVVGVT SWGYGCGVKD SPGVYTKVSAFVPW : : : . : . DSCQGDSSGGPLMC EKPDES WVVGVT SWGYGCGVKDTPGVYTRVPAFVPW	250
IKSVTKL . IKSVTSL	258

From the 258 amino acid sequence positions included in the comparison there are 233 amino acids that are identical in both compounds (upper sequence: compound of the formula I; lower sequence: compound of the formula II; identical amino acids are indicated by vertical lines).

The inventive neurotrypsins are unique when compared with the known serine proteases in that they are expressed according to currently available observations in a distinct degree in neurons. A further organ with a strong expression of neurotrypsin are the lungs (see Gschwend et al., Mol. Cell. Neurosci. 9, pages 207-219, 1997).

The proteins that are structurally most similar to the compounds of the formulas I or II are serine proteases, such as tissue-type plasminogen activator (tPA), urokinase-type plasminogen activator (uPA), plasmin, trypsin, apolipoprotein (a), coagulation factor XI, neuropsin, and acrosin.

5

In the adult brain, the inventive compounds are expressed predominantly in the cerebral cortex, the hippocampus, and the amygdala.

10 In the adult brain stem and the spinal cord, the inventive compounds are expressed predominantly in the motor neurons. A slightly weaker expression is found in the neurons of the superficial layers of the dorsal horn of the spinal cord.

In the adult peripheral nervous system, the inventive compounds are expressed in a subpopulation of the sensory ganglia neurons.

15

The inventive compounds were found in connection with a study aimed at discovering trypsin-like serine proteases in the nervous system.

20 The first compound that was found and characterized was the compound of the formula II (Gschwend et al., Mol. Cell. Neurosci. 9, pages 207-219, 1997).

25 By means of an alignment of the protease domains of 7 known serine proteases (tissue-type plasminogen activator, urokinase-type plasminogen activator, thrombin, plasmin, trypsin, chymotrypsin, and pancreatic elastase) in the proximity of the histidine and the serine of the catalytic triade of the active site, the sequences of the so-called primer oligonucleotides for the polymerase chain reaction were determined.

30 The primer oligonucleotides were used in a polymerase chain reaction (PCR) together with ss-cDNA from total RNA of the brains of 10 days old mice and resulted in the amplification of a cDNA fragment of a length of approximately 500 base pairs.

35 This cDNA fragment was used successfully for the isolation of further cDNA fragments by screening commercially available cDNA libraries. Together, the isolated cDNA fragments covered the full length of the coding part of the compound of the formula II.

By conventional DNA sequencing the complete nucleotide sequence and the amino acid sequence deduced therefrom was obtained.

5 The compound of the formula I was cloned based on its pronounced similarity with the compound of the formula II.

The primer oligonucleotides used were synthesized according to the known sequence of the compound of the formula II.

10

The cloning of the compound of the formula I was performed by means of two commercially available cDNA libraries from fetal human brain.

15 This procedure for the cloning can also be used for the isolation of the homologous compounds of other species, such as rat, rabbit, guinea pig, cow, sheep, pig, primates, birds, zebra fish (*Brachydanio rerio*), *Drosophila melanogaster*, *Caenorhabditis elegans* etc.

20 The coding nucleotide sequences can be used for the production of proteins with the coded amino acid sequences of the compounds of the formulas I or II. A procedure developed in our laboratory allows the production of recombinant proteins in myeloma cells as fusion proteins with an immunoglobulin domain (constant domain of the kappa light chain). The principle of the construction is given in detail by Rader et al. (Rader et al., Eur. J. Biochem. 215, pages 133-141, 1993). The fusion protein produced by the
25 myeloma cells was isolated by immunoaffinity chromatography using a monoclonal antibody against the Ig domain of the kappa light chain. With the same expression method, also the native protein of a compound, starting from the coding sequence, can be produced.

30 The coding sequences of the compounds of the formulas I or II can be used as starting compounds for the discovery and the isolation of alleles of the compounds of the formulas I or II. Both the polymerase chain reaction and the nucleic acid hybridization can be used for this purpose.

The coding sequences of the compounds of the formulas I or II can be used as starting compounds for so-called "site-directed mutagenesis", in order to generate nucleotide sequences coding the coded proteins that are defined by the compounds of the formulas I or II, or parts thereof, but whose nucleotide sequence is degenerated with respect to the compounds of the formulas I or II due to use of alternative codons.

5

The coding sequences of the compounds of the formulas I or II can be used as starting compounds for the production of sequence variants by means of so-called site-directed mutagenesis.

10

Best Modes for Carrying out the Invention (Examples)**cDNA cloning of the compound of the formula II (neurotrypsin of the mouse)**

- 5 Total RNA was isolated from the brains of 10 days old mice (ICR-ZUR) according to the method of Chomczynski and Sacchi (1987). The production of single stranded cDNA was carried out using oligo(dT) primer and a RNA-dependent DNA polymerase (SuperScript RNase H⁻Reverse Transcriptase; Gibco BRL, Gaithersburg, MD) according to the instruction of the supplier. For the realization of the polymerase chain reaction one
- 10 forward primer was synthesized based on the amino acid sequence of the region of the conserved histidine of the catalytic triade and one primer in the backward direction was synthesized based on the amino acid sequence of the region of the conserved serine of the catalytic triade of the serine proteases. The amino acid sequences used for the determination of the oligonucleotide primers were taken from seven known serine
- 15 proteases. They are presented in the following.

Protease domain	I →		← II	
	N	H	S	C
tPA (m)	..SSC	W V L S A A H C	FLE.....HDA	C Q G D S G G PLV..
uPA (m)	..SPC	W V A S A A H C	FIQ.....TDS	C K G D S G G PLI..
thrombin (m)	..SDR	W V L T A A H C	ILY.....GDA	C E G D S G G PFV..
plasmin (m)	..APE	W V L T A A H C	LKS.....VDS	C Q G D S G G PLV..
trypsin (m)	..NDQ	W V V S A A H C	YKY.....KDS	C Q G D S G G PVV..
chymTryp b (r)	..SED	W V V T A A H C	GVK.....VSS	C M G D S G G PLV..
pancElas II (m)	..ANN	W V L T A A H C	LSN.....TSS	C N G D S G G PLN..

Primer (I) 5'-TCG GTI SYI WSI GCI GCI CAT TG-3' (II) 3'-ACR BTY CCI CTR WSI CCI CC-5'

- The protease domains of 7 known serine proteases (tissue-type plasminogen activator, urokinase-type plasminogen activator, thrombin, plasmin, trypsin, chymotrypsin, and pancreatic elastase) were aligned in the region of the conserved
- 20 histidine and serine of the catalytic triade of the active site. The conserved amino acids of these regions were taken as the basis for the determination of the degenerated primers. The primer sequences are given according to the recommendation of the IUB nomenclature (Nomenclature Committee 1985).

- 25 The primers used in the PCR contained restriction sites for *EcoRI* and *BamHI* at their 5' ends in order to facilitate a subsequent cloning.

The following primers were used:

In the reading direction (sense primers):

5'-GGGGAATTCTGGGTI(C/G)(T/C)I(T/A)(G/C)IGCIGCICA(T/C)TG-3'

5 In the counter direction (antisense primers):

5'-GGGGGATCCCCICCI(G/C)(A/T)(A/G)TCICC(C/T)T(G/C/T)(G/A)CA-3'.

10 The polymerase chain reaction was carried out under standard conditions using the DNA polymerase AmpliTaq (Perkin Elmer) according to the recommendations of the producer. The following PCR profile was employed: 93°C for 3 minutes, followed by 35 cycles of 93°C for 1 minute, 48°C for 2 minutes, and 72°C for 2 minutes. Following the last cycle, the incubation was continued at 72°C for further 10 minutes.

15 The amplified fragments had an approximate length of 500 base pairs. They were cut with *EcoRI* and *BamHI* and inserted in a Blue Script vector (Bluescript SK(-), Stratagene). The resulting clones were analyzed by DNA sequence determination using the dideoxy chain termination method (Sanger et al., Proc. Natl. Acad. Sci. USA 77, pages 2163-2167, 1977) on an automated DNA sequencer (LI-COR, model 4000L; Lincoln, NE) using a commercial sequencing kit (SequiTerm long-read cycle sequencing
20 kit-LC; Epicentre Technologies, Madison, WI). The analysis yielded a sequence of 474 base pairs of the catalytic region of the serine protease domain of the compound of the formula II.

25 The 474 base pair long PCR fragment was used for screening of an oligo(dT)-primed Uni-ZAP-XR cDNA library from the brain of 20 days old mice (Stratagene; cat. no. 937 319). At total of 3×10^8 lambda plaques were screened under high stringent conditions (Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, 1989) using a radioactively labeled PCR fragment as a probe and 24 positive clones were found.

30

From the positive Lambda-Uni-ZAP-XR phagemid clones the corresponding Bluescript plasmid was cut out by *in vivo* excision according to a standard method recommended by the producer (Stratagene). In order to determine the length of the inserted fragments the corresponding Bluescript plasmid clones were digested with *SacI*
35 and *KpnI*. The clones containing the longest fragments were analyzed by DNA

sequencing (as described above) and for subsequent data analysis the GCG software (version 8.1, Unix; Silicon Graphics, Inc.) was used.

Because none of the clones contained the coding sequence in full length, a second
5 cDNA library was screened. The library used in this screen was an oligo(dT)- and
random-primed cDNA library in a Lambda phage (Lambda gt10) which was based on
mRNA from 15 days old mouse embryos (oligo(dT)- and random-primed Lambda gt10
cDNA library; Clontech, Palo Alto, CA; cat. no. ML 3002a). As a probe a radioactively
10 labeled DNA fragment (Aval/AatII) from the 5' end of the longest clone of the first screen
was used and approximately 2×10^8 plaques were screened. This screen resulted in 14
positive clones. The cDNA fragments were excised with *EcoRI* and cloned into the
Bluecript vector (KS(+); Stratagene). The sequence analysis was carried out as
described above.

15 In this way the nucleotide sequence over the full length cDNA of 2361 and 2376
base pairs, respectively, of the compound of the formula II was obtained. With the
described procedure of PCR cloning it is possible to find and isolate also variant forms of
the compounds of the formulas I or II, as for example their alleles or their splice variants.
The described method of screening of a cDNA library allows also the detection and the
20 isolation of compounds which hybridize under stringent conditions with the coding
sequences of the compounds of the formulas I or II.

Cloning of the cDNA of the compound of the formula I (neurotrypsin of the human)

The cloning of the cDNA of the compound of the formula I was carried out basing
5 on the nucleotide sequence of the compound of the formula II. As a first step, a fragment
of the compound of the formula I was amplified using the polymerase chain reaction
(PCR). As a matrix we used the DNA obtained from a cDNA library from the brain of a
human fetus (17th - 18th week of pregnancy) which is commercially available (Oligo(dT)-
and random-primed, human fetal brain cDNA library in the Lambda ZAP II vector, cat.
10 no. 936206, Stratagene). The synthetic PCR primers contained restriction sites for
HindIII and *XhoI* at the 5' end in order to facilitate the subsequent cloning.

In the reading direction (sense primers):

5'-GGGAAGCTTGGICA(A/G)TGGGGIACI(A/G)TITG(C/T)GA(C/T)-3'

15 In the counter direction (antisense primers):

5'-GGGCTCGAGCCCCAICCTGTTATGTAAIAGTTG-3'

The PCR was carried out under standard conditions using the DNA polymerase
20 Amplitaq (Perkin Elmer) according to the recommendations of the producer. The
resulting fragment of 1116 base pairs was inserted into the Bluescript vector (Bluescript
SK(-), Stratagene). A 600 base pairs long *HindIII/StuI* fragment, corresponding to the 5'
half the 1116 base pairs long PCR fragment, was used for the screening of a Lambda
cDNA library from human fetal brain (Human Fetal Brain 5'-STRETCH PLUS cDNA
25 library; Lambda gt10; cat. no. HL 3003 a; Clontech). 2x10⁶ Lambda plaques were
screened under high stringent conditions (Sambrook et al., Molecular Cloning: A
laboratory manual, Cold Spring Harbor Laboratory Press, 1989) by means of a
radioactively labeled PCR fragment, and 23 positive clones were found and isolated.

30 From the positive Lambda gt10 clones the corresponding cDNA fragments were
excised with *EcoRI* and inserted into a Bluescript vector (Bluescript KS(+), Stratagene).
The sequencing was carried out by means of the dideoxy chain termination method
(Sanger et al., Proc. Natl. Acad. Sci. USA 77, pages 2163-2167, 1977), using a
commercial sequencing kit (SequiTherm long-read cycle sequencing kit-LC; Epicentre
35 Technologies, Madison, WI) and Bluescript-specific primers.

In an alternative sequencing strategy, the cDNA fragments of the positive Lambda gt10 clones were PCR amplified using Lambda-specific primers. The sequencing was carried out as described above.

5

The computerized analysis of the sequences was performed by means of the program package GCG (version 8.1, Unix; Silicon Graphics Inc.).

In this way the nucleotide sequence over the full length of the cDNA of 3350 base pairs was obtained. With the described procedure for PCR cloning it is possible to find and to isolate also variant forms of the compounds of the formulas I or II, as for example their alleles or their splice variants. The described procedure for the screening of a cDNA library allows also the discovery and the isolation of compounds which hybridize under stringent conditions with the coding sequences of the compounds of the formulas I or II.

15

Visualization of the coded sequences of the compounds of the formulas I or II by means of antibodies

5 The more than 60 amino acids long proline-rich, basic segment at the amino terminus of the coded sequence of the compounds of the formulas I or II is well suited for the production of antibodies by means of synthesizing peptides and using them for immunization. We have selected two peptide sequences with a length of 19 and 13 amino acids from the proline-rich, basic segment at the amino terminus of the coded
10 sequence of the compound of the formula II for the generation of antibodies. The peptides had the following sequences:

Peptide 1: $\text{H}_2\text{N-SRS PLH RPH PSP PRS QX-CONH}_2$

Peptide 2: $\text{H}_2\text{N-LPS SRR PPR TPR F-COOH}$

15 The two peptides were synthesized chemically, coupled to a macromolecular carrier (Keyhole Limpet Hemacyanin), and injected into 2 rabbits for immunization. The resulting antisera exhibit a high antibody titer and could successfully be used both for the identification of native neurotrypsin in brain extract of the mouse and for the identification of recombinant neurotrypsin. The employed procedure for the generation of antibodies
20 can also be used for the generation of antibodies against the coded sequence of the compound of the formula I.

25 The resulting antibodies against the partial sequences of the coded sequences of the compounds of the formulas I or II can be used for the detection and the isolation of variant forms of the compounds of the formulas I or II, as for example alleles or splice variants. Such antibodies can also be used for the detection and isolation of gene technologically generated variants of the compounds of the formulas I or II.

Purification of the coded sequences of the compounds of the formulas I or II

Besides conventional chromatographic methods, as for example ion exchange chromatography, the purification of the coded sequences of the compounds of the formulas I or II can also be achieved using two affinity chromatographic purification procedures. One affinity chromatographic purification procedure is based on the availability of antibodies. By coupling the antibodies on a chromatographic matrix, a purification procedure results, in which a very high degree of purity of the corresponding compound can be achieved in one step.

Another important feature that can be used for the purification of the coded sequences of the compounds of the formulas I or II is the proline-rich, basic segment at the amino terminus. It may be expected that, due to the high density of positive charges, this segment mediates the binding of the coded sequences of the compounds of the formulas I or II to heparin and heparin-like affinity matrices. This principle allows also the isolation, or at least the enrichment, of variant forms of the coded sequences of the compounds of the formulas I or II, as for example their alleles or splice variants. Likewise the heparin affinity chromatography can be used for the isolation, or at least the enrichment, of species-homologous proteins of the compounds of the formulas I or II.

Industrial Applicability

The coding sequences of the formulas I and II can be used for the production of the coded proteins or parts thereof of the formulas I and II. The production of the coded
5 proteins can be achieved in procaryotic or eucaryotic expression systems.

The gene expression pattern of the inventive compounds in the brain is extremely interesting, because these molecules are expressed in the adult nervous system predominantly in neurons of those regions that are thought to play an important role in
10 learning and memory functions. Together with the recently found evidence for a role of extracellular proteases in neural plasticity, the expression pattern allows the assumption that the proteolytic activity of neurotrypsin has a role in structural reorganizations in connection with learning and memory operations, for example operations which are
15 involved in the processing and storage of learned behaviors, learned emotions, or memory contents. The inventive compounds may, thus, represent a target for pharmaceutical intervention in malfunctions of the brain.

The gene expression pattern of the inventive compounds in the cerebral cortex (especially layers V and VI) is extremely interesting, because a reduction of the cellular
20 differentiation in the cerebral cortex has been found to be associated with schizophrenia. The inventive compounds may, thus, be a target for pharmaceutical intervention in schizophrenia and related psychiatric diseases.

The coding sequences of the inventive compounds have been found to be
25 increased in the neurons located adjacent to the damaged tissue of a focal ischemic stroke, indicating that the inventive compounds play a role in the tissue reaction in the injured cerebral tissue. The inventive compounds may, thus, represent a target for pharmaceutical intervention after ischemic stroke and other forms of neural tissue
30 damage.

Tissue-type plasminogen activator, a serine protease related to the inventive compounds, has recently been found to be involved in excitotoxicity-mediated neuronal cell death. A similar function is conceivable for the inventive compounds and, thus, the
35 inventive compounds represent a possible target for a pharmacological intervention in diseases in which cell death occurs.

The gene expression pattern of the inventive compounds in the spinal cord and in the sensory ganglia is interesting, because these molecules are expressed in the adult nervous system in neurons of those brain regions that are thought to play a role in the processing of pain, as well as in the pathogenesis of pathological pain. The inventive compounds may, thus, be a target for pharmaceutical intervention in pathological pain.

10 In the following part statements concerning the compounds of the formulas I or II are given:

(1) INFORMATION ABOUT THE COMPOUND OF THE FORMULA I(Neurotrypsin of the human)

(i) SEQUENCE CHARACTERISTICS:

5

- (A) LENGTH: 3350 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single strand
- (D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: cDNA to mRNA

(vi) ORIGINAL SOURCE:

15

- (A) ORGANISM: Homo sapiens
- (D) DEVELOPMENT STAGE: fetal
- (F) TISSUE TYPE: brain

(vii) IMMEDIATE SOURCE:

20

- (A) LIBRARY: human fetal brain 5'-stretch plus cDNA library in the lambda gt10 vector; catalog No. HL 3003a; Clontech, Palo Alto, CA, USA.

(B) CLONE: cDNA Clone No.:

25

3-1, 3-2, 3-6, 3-7, 3-8, 3-10, 3-11, 3-12

(ix) FEATURE:

30

- (A) NAME/KEY: Signal peptide
- (B) LOCATION: 44 .. 103

(ix) FEATURE:

(A) NAME/KEY: mature peptide

(B) LOCATION: 104 .. 2668

5

(ix) FEATURE:

(A) NAME/KEY: coding sequence

10 (B) LOCATION: 44 .. 2668

(ix) FEATURE:

15 (A) NAME/KEY: Proline-rich, basic segment

(B) LOCATION: 104 .. 319

(ix) FEATURE:

20

(A) NAME/KEY: Kringle domain

(B) LOCATION: 320 .. 538

25 (ix) FEATURE:

(A) NAME/KEY: SRCR domain 1

(B) LOCATION: 551 .. 856

30

(ix) FEATURE:

(A) NAME/KEY: SRCR domain 2

(B) LOCATION: 881 .. 1186

35

(ix) FEATURE:

(A) NAME/KEY: SRCR domain 3

5 (B) LOCATION: 1202 .. 1504

(ix) FEATURE:

10 (A) NAME/KEY: SRCR domain 4

(B) LOCATION: 1541 .. 1846

(ix) FEATURE:

15

(A) NAME/KEY: proteolytic domain

(B) LOCATION: 1898 .. 2668

20 (ix) FEATURE:

(A) NAME/KEY: histidine of the catalytic triade

(B) LOCATION: 2069 - 2071

25

(ix) FEATURE:

(A) NAME/KEY: aspartic acid of the catalytic triade

(B) LOCATION: 2219 - 2221

30

(ix) FEATURE:

(A) NAME/KEY: serine of the catalytic triade

35 (B) LOCATION: 2516 .. 2518

(ix) FEATURE:

- 5 (A) NAME/KEY: polyA signal
(B) LOCATION: 2873 .. 2878

(ix) FEATURE

10

- (A) NAME/KEY: polyA signal
(B) LOCATION: 3034 .. 3039

15 (ix) FEATURE:

- (A) NAME/KEY: polyA signal
(B) LOCATION: 3215 .. 3220

20

(ix) FEATURE:

- (A) NAME/KEY: 3'UTR
(B) LOCATION: 2669 .. 3350

25

(ix) FEATURE

- (A) NAME/KEY: 5'UTR
30 (B) LOCATION: 1 .. 43

- 20 -

Comparison of the formula I (neur trypsin of the human)

CGGAAGCTGG	GGAGCATGGA	CCAGACCCCG	CAGCGCTGGC	ACC	ATG	ACG	CTC	GCC	55
					Met	Thr	Leu	Ala	
					-20				
CGC	TTC	GTG	CTA	GCC	CTG	ATG	TTA	GGG	103
Arg	Phe	Val	Leu	Ala	Leu	Met	Leu	Gly	
-15					-10			-5	
									-1
TTT	GAT	TCT	GTC	CTC	AAT	GAT	TCC	CTC	151
Phe	Asp	Ser	Val	Leu	Asn	Asp	Ser	Leu	
1				5				10	
									15
CCC	CCT	GCG	GGT	CCG	CAC	TAC	CCC	TAT	199
Pro	Pro	Ala	Gly	Pro	His	Tyr	Pro	Tyr	
			20					25	
									30
CCC	CCG	ACG	ACG	CGT	CCG	CCG	CCG	CCT	247
Pro	Pro	Thr	Thr	Arg	Pro	Pro	Pro	Leu	
			35					40	
									45
CCG	CGG	GCG	CTC	CCT	GCC	CAG	CGC	CCG	295
Pro	Arg	Ala	Leu	Pro	Ala	Gln	Arg	Pro	
			50					55	
									60
ACG	CCC	CGG	CCG	CAC	CCC	TGG	GGC	TGC	343
Thr	Pro	Arg	Pro	His	Pro	Trp	Gly	Cys	
65					70			75	
									80
AGC	GTG	ACG	GAC	TTC	GGC	GCC	CCG	TGT	391
Ser	Val	Thr	Asp	Phe	Gly	Ala	Pro	Cys	
				85				90	
									95
CCC	TTC	CTG	GAG	CGG	TCG	CCC	CCA	GCG	439
Pro	Phe	Leu	Glu	Arg	Ser	Pro	Pro	Ala	
			100					105	
									110
CAG	CGC	CAC	AAC	TTT	TGT	CGG	AGC	CCC	487
Gln	Arg	His	Asn	Phe	Cys	Arg	Ser	Pro	
			115					120	
									125
TGT	TTC	TAC	GGA	GAC	GCC	CGT	GGC	AAG	535
Cys	Phe	Tyr	Gly	Asp	Ala	Arg	Gly	Lys	
			130					135	
									140
TGC	AGA	CAC	GGA	TCA	GTA	CGA	CTT	CGT	583
Cys	Arg	His	Gly	Ser	Val	Arg	Leu	Arg	
145					150			155	
									160
GGC	ACA	GTG	GAA	GTA	TAT	GCA	AGT	GGA	631
Gly	Thr	Val	Glu	Val	Tyr	Ala	Ser	Gly	
				165				170	
									175
AGC	CAC	TGG	GAT	GAT	TCT	GAT	GCA	TCA	679
Ser	His	Trp	Asp	Asp	Ser	Asp	Ala	Ser	
				180				185	
									190

CTG GGA GGA AAA GGA ATA GCA AAA CAA ACC CCG TTT TCT GGA CTG GGC Leu Gly Gly Lys Gly Ile Ala Lys Gln Thr Pro Phe Ser Gly Leu Gly 195 200 205	727
CTT ATT CCC ATT TAT TGG AGC AAT GTC CGT TGC CGA GGA GAT GAA GAA Leu Ile Pro Ile Tyr Trp Ser Asn Val Arg Cys Arg Gly Asp Glu Glu 210 215 220	775
AAT ATA CTG CTT TGT GAA AAA GAC ATC TGG CAG GGT GGG GTG TGT CCT Asn Ile Leu Leu Cys Glu Lys Asp Ile Trp Gln Gly Gly Val Cys Pro 225 230 235 240	823
CAG AAG ATG GCA GCT GCT GTC ACG TGT AGC TTT TCC CAT GGC CCA ACG Gln Lys Met Ala Ala Val Thr Cys Ser Phe Ser His Gly Pro Thr 245 250 255	871
TTC CCC ATC ATT CGC CTT GCT GGA GGC AGC AGT GTG CAT GAA GGC CGG Phe Pro Ile Ile Arg Leu Ala Gly Gly Ser Ser Val His Glu Gly Arg 260 265 270	919
GTG GAG CTC TAC CAT GCT GGC CAG TGG GGA ACC GTT TGT GAT GAC CAA Val Glu Leu Tyr His Ala Gly Gln Trp Gly Thr Val Cys Asp Asp Gln 275 280 285	967
TGG GAT GAT GCC GAT GCA GAA GTG ATC TGC AGG CAG CTG GGC CTC AGT Trp Asp Asp Ala Asp Ala Val Ile Cys Arg Gln Leu Gly Leu Ser 290 295 300	1015
GGC ATT GCC AAA GCA TGG CAT CAG GCA TAT TTT GGG GAA GGG TCT GGC Gly Ile Ala Lys Ala Trp His Gln Ala Tyr Phe Gly Glu Gly Ser Gly 305 310 315 320	1063
CCA GTT ATG TTG GAT GAA GTA CGC TGC ACT GGG AAT GAG CTT TCA ATT Pro Val Met Leu Asp Glu Val Arg Cys Thr Gly Asn Glu Leu Ser Ile 325 330 335	1111
GAG CAG TGT CCA AAG AGC TCC TGG GGA GAG CAT AAC TGT GGC CAT AAA Glu Gln Cys Pro Lys Ser Ser Trp Gly Glu His Asn Cys Gly His Lys 340 345 350	1159
GAA GAT GCT GGA GTG TCC TGT ACC CCT CTA ACA GAT GGG GTC ATC AGA Glu Asp Ala Gly Val Ser Cys Thr Pro Leu Thr Asp Gly Val Ile Arg 355 360 365	1207
CTT GCA GGT GGG AAA GGC AGC CAT GAG GGT CGC TTG GAG GTA TAT TAC Leu Ala Gly Gly Lys Gly Ser His Glu Gly Arg Leu Glu Val Tyr Tyr 370 375 380	1255
AGA GGC CAG TGG GGA ACT GTC TGT GAT GAT GGC TGG ACT GAG CTG AAT Arg Gly Gln Trp Gly Thr Val Cys Asp Asp Gly Trp Thr Glu Leu Asn 385 390 395 400	1303
ACA TAC GTG GTT TGT CGA CAG TTG GGA TTT AAA TAT GGT AAA CAA GCA Thr Tyr Val Val Cys Arg Gln Leu Gly Phe Lys Tyr Gly Lys Gln Ala 405 410 415	1351
TCT GCC AAC CAT TTT GAA GAA AGC ACA GGG CCC ATA TGG TTG GAT GAC Ser Ala Asn His Phe Glu Glu Ser Thr Gly Pro Ile Trp Leu Asp Asp 420 425 430	1399

- 22 -

GTC	AGC	TGC	TCA	GGA	AAG	GAA	ACC	AGA	TTT	CTT	CAG	TGT	TCC	AGG	CGA	1447
Val	Ser	Cys	Ser	Gly	Lys	Glu	Thr	Arg	Phe	Leu	Gln	Cys	Ser	Arg	Arg	
		435					440					445				
CAG	TGG	GGA	AGG	CAT	GAC	TGC	AGC	CAC	CGC	GAA	GAT	GTT	AGC	ATT	GCC	1495
Gln	Trp	Gly	Arg	His	Asp	Cys	Ser	His	Arg	Glu	Asp	Val	Ser	Ile	Ala	
	450					455					460					
TGC	TAC	CCT	GGC	GGC	GAG	GGA	CAC	AGG	CTC	TCT	CTG	GGT	TTT	CCT	GTC	1543
Cys	Tyr	Pro	Gly	Gly	Glu	Gly	His	Arg	Leu	Ser	Leu	Gly	Phe	Pro	Val	
465					470					475					480	
AGA	CTG	ATG	GAT	GGA	GAA	AAT	AAG	AAA	GAA	GGA	CGA	GTG	GAG	GTT	TTT	1591
Arg	Leu	Met	Asp	Gly	Glu	Asn	Lys	Lys	Glu	Gly	Arg	Val	Glu	Val	Phe	
				485					490					495		
ATC	AAT	GGC	CAG	TGG	GGA	ACA	ATC	TGT	GAT	GAT	GGA	TGG	ACT	GAT	AAG	1639
Ile	Asn	Gly	Gln	Trp	Gly	Thr	Ile	Cys	Asp	Asp	Gly	Trp	Thr	Asp	Lys	
			500					505					510			
GAT	GCA	GCT	GTG	ATC	TGT	CGT	CAG	CTT	GGC	TAC	AAG	GGT	CCT	GCC	AGA	1687
Asp	Ala	Ala	Val	Ile	Cys	Arg	Gln	Leu	Gly	Tyr	Lys	Gly	Pro	Ala	Arg	
		515					520					525				
GCA	AGA	ACC	ATG	GCT	TAC	TTT	GGA	GAA	GGA	AAA	GGA	CCC	ATC	CAT	GTG	1735
Ala	Arg	Thr	Met	Ala	Tyr	Phe	Gly	Glu	Gly	Lys	Gly	Pro	Ile	His	Val	
	530					535					540					
GAT	AAT	GTG	AAG	TGC	ACA	GGA	AAT	GAG	AGG	TCC	TTG	GCT	GAC	TGT	ATC	1783
Asp	Asn	Val	Lys	Cys	Thr	Gly	Asn	Glu	Arg	Ser	Leu	Ala	Asp	Cys	Ile	
545					550					555					560	
AAG	CAA	GAT	ATT	GGA	AGA	CAC	AAC	TGC	CGC	CAC	AGT	GAA	GAT	GCA	GGA	1831
Lys	Gln	Asp	Ile	Gly	Arg	His	Asn	Cys	Arg	His	Ser	Glu	Asp	Ala	Gly	
				565					570					575		
GTT	ATT	TGT	GAT	TAT	TTT	GGC	AAG	AAG	GCC	TCA	GGT	AAC	AGT	AAT	AAA	1879
Val	Ile	Cys	Asp	Tyr	Phe	Gly	Lys	Lys	Ala	Ser	Gly	Asn	Ser	Asn	Lys	
			580					585					590			
GAG	TCC	CTC	TCA	TCT	GTT	TGT	GGC	TTG	AGA	TTA	CTG	CAC	CGT	CGG	CAG	1927
Glu	Ser	Leu	Ser	Ser	Val	Cys	Gly	Leu	Arg	Leu	Leu	His	Arg	Arg	Gln	
		595					600					605				
AAG	CGG	ATC	ATT	GGT	GGG	AAA	AAT	TCT	TTA	AGG	GGT	GGT	TGG	CCT	TGG	1975
Lys	Arg	Ile	Ile	Gly	Gly	Lys	Asn	Ser	Leu	Arg	Gly	Gly	Trp	Pro	Trp	
	610					615					620					
CAG	GTT	TCC	CTC	CGG	CTG	AAG	TCA	TCC	CAT	GGA	GAT	GGC	AGG	CTC	CTC	2023
Gln	Val	Ser	Leu	Arg	Leu	Lys	Ser	Ser	His	Gly	Asp	Gly	Arg	Leu	Leu	
625					630					635					640	
TGC	GGG	GCT	ACG	CTC	CTG	AGT	AGC	TGC	TGG	GTC	CTC	ACA	GCA	GCA	CAC	2071
Cys	Gly	Ala	Thr	Leu	Leu	Ser	Ser	Cys	Trp	Val	Leu	Thr	Ala	Ala	His	
				645					650					655		
TGT	TTC	AAG	AGG	TAT	GGC	AAC	AGC	ACT	AGG	AGC	TAT	GCT	GTT	AGG	GTT	2119
Cys	Phe	Lys	Arg	Tyr	Gly	Asn	Ser	Thr	Arg	Ser	Tyr	Ala	Val	Arg	Val	
			660					665					670			

GGA GAT TAT CAT ACT CTG GTA CCA GAG GAG TTT GAG GAA GAA ATT GGA Gly Asp Tyr His Thr Leu Val Pro Glu Glu Phe Glu Glu Ile Gly 675 680 685	2167
GTT CAA CAG ATT GTG ATT CAT CGG GAG TAT CGA CCC GAC CGC AGT GAT Val Gln Gln Ile Val Ile His Arg Glu Tyr Arg Pro Asp Arg Ser Asp 690 695 700	2215
TAT GAC ATA GCC CTG GTT AGA TTA CAA GGA CCA GAA GAG CAA TGT GCC Tyr Asp Ile Ala Leu Val Arg Leu Gln Gly Pro Glu Glu Gln Cys Ala 705 710 715 720	2263
AGA TTC AGC AGC CAT GTT TTG CCA GCC TGT TTA CCA CTC TGG AGA GAG Arg Phe Ser Ser His Val Leu Pro Ala Cys Leu Pro Leu Trp Arg Glu 725 730 735	2311
AGG CCA CAG AAA ACA GCA TCC AAC TGT TAC ATA ACA GGA TGG GGT GAC Arg Pro Gln Lys Thr Ala Ser Asn Cys Tyr Ile Thr Gly Trp Gly Asp 740 745 750	2359
ACA GGA CGA GCC TAT TCA AGA ACA CTA CAA CAA GCA GCC ATT CCC TTA Thr Gly Arg Ala Tyr Ser Arg Thr Leu Gln Gln Ala Ala Ile Pro Leu 755 760 765	2407
CTT CCT AAA AGG TTT TGT GAA GAA CGT TAT AAG GGT CGG TTT ACA GGG Leu Pro Lys Arg Phe Cys Glu Glu Arg Tyr Lys Gly Arg Phe Thr Gly 770 775 780	2455
AGA ATG CTT TGT GCT GGA AAC CTC CAT GAA CAC AAA CGC GTG GAC AGC Arg Met Leu Cys Ala Gly Asn Leu His Glu His Lys Arg Val Asp Ser 785 790 795 800	2503
TGC CAG GGA GAC AGC GGA GGA CCA CTC ATG TGT GAA CGG CCC GGA GAG Cys Gln Gly Asp Ser Gly Gly Pro Leu Met Cys Glu Arg Pro Gly Glu 805 810 815	2551
AGC TGG GTG GTG TAT GGG GTG ACC TCC TGG GGG TAT GGC TGT GGA GTC Ser Trp Val Val Tyr Gly Val Thr Ser Trp Gly Tyr Gly Cys Gly Val 820 825 830	2599
AAG GAT TCT CCT GGT GTT TAT ACC AAA GTC TCA GCC TTT GTA CCT TGG Lys Asp Ser Pro Gly Val Tyr Thr Lys Val Ser Ala Phe Val Pro Trp 835 840 845	2647
ATA AAA AGT GTC ACC AAA CTG TAA TTCTTCATGG AAACCTCAAA GCAGCATTT Ile Lys Ser Val Thr Lys Leu * 850 855	2700
AAACAAATGG AAAACTTTGA ACCCCCACTA TTAGCACTCA GCAGAGATGA CAACAAATGG	2760
CAAGATCTGT TTTTGCTTTG TGTTGTGGTA AAAAATTGTG TACCCCTGC TGCTTTTGAG	2820
AAATTTGTGA ACATTTTCAG AGGCCTCAGT GTAGTGAAG TGATAATCCT TAAATGAACA	2880
TTTTCTACCC TAATTTCACT GGAGTGACTT ATTCTAAGCC TCATCTATCC CCTACCTATT	2940

TCTCAAAATC ATTCTATGCT GATTTTACAA AAGATCATTT TTACATTTGA ACTGAGAACC 3000
CCTTTTAATT GAATCAGTGG TGTCTGAAAT CATATTAAAT ACCCACATTT GACATAAATG 3060
CGGTACCCTT TACTACACTC ATGAGTGGCA TATTTATGCT TAGGTCTTTT CAAAAGACTT 3120
GACAAGAAAT CTTCATATTC TCTGTAGCCT TTGTCAAGTG AGGAAATCAG TGGTTAAAGA 3180
ATTCCACTAT AAACTTTTAG GCCTGAATAG GAGTAGTAAA GCCTCAAGGA CATCTGCCTG 3240
TCACAATATA TTCTCAAAGT GATCTGATAT TTGGAAACAA GTATCCTTGT TGAGTACCAA 3300
GTGCTACAGA AACCATAAGA TAAAAATACT TTCTACCTAC AGCGTGCCCG 3350

(1) INFORMATION ABOUT THE COMPOUND OF THE FORMULA II (Neurotrypsin of the mouse)

(i) SEQUENCE CHARACTERISTICS:

5

- (A) LENGTH: 2376 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single strand
- (D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: cDNA to mRNA

(vi) ORIGINAL SOURCE:

15

- (A) ORGANISM: Mus musculus
- (D) DEVELOPMENT STAGE: postnatal day 10
- (F) TISSUE TYPE: brain
- (G) CELL TYPE: neurons

20

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: mouse brain cDNA library in the lambda Uni-ZAP-XR vector, oligo(dT)-primed, from Balb c mice, postnatal day 20, Cat. No.. 937 319; Stratagene, La Jolla, CA, USA

25

- (B) CLONE: cDNA clone no. 16

(vii) IMMEDIATE SOURCE:

30

- (A) LIBRARY: mouse brain cDNA library in the Lambda gt10 vector, oligo(dT)- and random-primed, embryonic day 15, Cat. No. ML 3002a; Clontech, Palo Alto, CA, USA

35

- (B) CLONE: cDNA clone #25

(ix) FEATURE:

(A) NAME/KEY: signal peptide

5 (B) LOCATION: 24 .. 86

(ix) FEATURE:

10 (A) NAME/KEY: mature peptide

(B) LOCATION: 87 .. 2306

(ix) FEATURE:

15

(A) NAME/KEY: coding sequence

(B) LOCATION: 24 .. 2306

20 (ix) FEATURE:

(A) NAME/KEY: proline-rich, basic segment

(B) LOCATION: 90 .. 275

25

(ix) FEATURE:

(A) NAME/KEY: Kringle domain

(B) LOCATION: 276 .. 494

30

(ix) FEATURE:

(A) NAME/KEY: SRCR domain 1

35 (B) LOCATION: 519 .. 824

(ix) FEATURE:

- 5 (A) NAME/KEY: SRCR domain 2
(B) LOCATION: 840 .. 1142

(ix) FEATURE:

10

- (A) NAME/KEY: SRCR domain 3
(B) LOCATION: 1179 .. 1484

15 (ix) FEATURE:

- (A) NAME/KEY: proteolytic domain
(B) LOCATION: 1536 .. 2306

20

(ix) FEATURE:

- (A) NAME/KEY: histidine of the catalytic triade
(B) LOCATION: 1707 .. 1709

25

(ix) FEATURE:

- (A) NAME/KEY: aspartic acid of the catalytic triade
30 (B) LOCATION: 1857 .. 1859

(ix) FEATURE:

- 35 (A) NAME/KEY: serine of the catalytic triade

- 28 -

(B) LOCATION: 2154 .. 2156

(ix) FEATURE:

5 (A) NAME/KEY: polyA signal

(B) LOCATION: 2324 .. 2329 and 2331 .. 2336

(ix) FEATURE:

10 (A) NAME/KEY: polyA segment

(B) LOCATION: 2357 .. 2376

(ix) FEATURE:

15

(A) NAME/KEY: 3'UTR

(B) LOCATION: 2307 .. 2341 or 2307 .. 2356

20

(ix) FEATURE:

(A) NAME/KEY: 5'UTR

(B) LOCATION: 1 .. 23

Coump und of the formula II (n ur trypsin of the mouse)

GGACCACACT CGGCGCCGCA GCC ATG GCG CTC GCC CGC TGC GTG CTG GCT GTG	53
Met Ala Leu Ala Arg Cys Val Leu Ala Val	
-20 -15	
ATT TTA GGG GCA CTG TCT GTA GTG GCC CGC GCT GAT CCG GTC TCG CGC	101
Ile Leu Gly Ala Leu Ser Val Val Ala Arg Ala Asp Pro Val Ser Arg	
-10 -5 1 5	
TCT CCC CTT CAC CGC CCG CAT CCG TCC CCA CCG CGT TCC CAA CAC GCG	149
Ser Pro Leu His Arg Pro His Pro Ser Pro Pro Arg Ser Gln His Ala	
10 15 20	
CAC TAC CTT CCC AGC TCG CGG CGG CCA CCC AGG ACC CCG CGC TTC CCG	197
His Tyr Leu Pro Ser Ser Arg Arg Pro Pro Arg Thr Pro Arg Phe Pro	
25 30 35	
CTC CCG CTG CGG ATC CCC GCT GCC CAG CGC CCG CAG GTC CTC AGC ACC	245
Leu Pro Leu Arg Ile Pro Ala Ala Gln Arg Pro Gln Val Leu Ser Thr	
40 45 50	
GGG CAC ACG CCC CCG ACG ATT CCA CGC CGC TGC GGG GCA GGA GAG TCG	293
Gly His Thr Pro Pro Thr Ile Pro Arg Arg Cys Gly Ala Gly Glu Ser	
55 60 65	
TGG GGC AAT GCC ACC AAC CTC GGC GTC CCG TGT CTA CAC TGG GAC GAG	341
Trp Gly Asn Ala Thr Asn Leu Gly Val Pro Cys Leu His Trp Asp Glu	
70 75 80 85	
GTG CCG CCC TTC CTG GAG CGG TCG CCC CCG GCC AGT TGG GCT GAG CTG	389
Val Pro Pro Phe Leu Glu Arg Ser Pro Pro Ala Ser Trp Ala Glu Leu	
90 95 100	
CGA GGG CAG CCG CAC AAC TTC TGC CGG AGC CCG GAT GGC TCG GGC AGA	437
Arg Gly Gln Pro His Asn Phe Cys Arg Ser Pro Asp Gly Ser Gly Arg	
105 110 115	
CCT TGG TGC TTC TAT CGG AAT GCC CAG GGC AAA GTA GAC TGG GGC TAC	485
Pro Trp Cys Phe Tyr Arg Asn Ala Gln Gly Lys Val Asp Trp Gly Tyr	
120 125 130	
TGC GAT TGT GGT CAA GGC CCG GCG TTG CCC GTC ATT CGC CTT GTT GGT	533
Cys Asp Cys Gly Gln Gly Pro Ala Leu Pro Val Ile Arg Leu Val Gly	
135 140 145	
GGG AAC AGT GGG CAT GAA GGT CGA GTG GAG CTG TAC CAC GCT GGC CAG	581
Gly Asn Ser Gly His Glu Gly Arg Val Glu Leu Tyr His Ala Gly Gln	
150 155 160 165	
TGG GGG ACC ATC TGT GAC GAC CAA TGG GAC AAT GCA GAC GCA GAC GTC	629
Trp Gly Thr Ile Cys Asp Asp Gln Trp Asp Asn Ala Asp Ala Asp Val	
170 175 180	
ATC TGT AGG CAG CTG GGG CTC AGT GGC ATT GCC AAA GCA TGG CAT CAG	677
Ile Cys Arg Gln Leu Gly Leu Ser Gly Ile Ala Lys Ala Trp His Gln	
185 190 195	

GCA CAT TTT GGG GAA GGA TCT GGC CCA ATA TTG TTG GAT GAA GTA CGC	725
Ala His Phe Gly Glu Gly Ser Gly Pro Ile Leu Leu Asp Glu Val Arg	
200 205 210	
TGC ACC GGA AAC GAG CTG TCA ATT GAG CAA TGT CCA AAG AGT TCC TGG	773
Cys Thr Gly Asn Glu Leu Ser Ile Glu Gln Cys Pro Lys Ser Ser Trp	
215 220 225	
GGC GAA CAT AAC TGT GGC CAT AAA GAA GAT GCT GGA GTG TCT TGT GTT	821
Gly Glu His Asn Cys Gly His Lys Glu Asp Ala Gly Val Ser Cys Val	
230 235 240 245	
CCT CTA ACA GAT GGT GTC ATC AGA CTG GCA GGA GGA AAA AGT ACC CAT	869
Pro Leu Thr Asp Gly Val Ile Arg Leu Ala Gly Gly Lys Ser Thr His	
250 255 260	
GAA GGT CGC CTG GAG GTC TAC TAC AAG GGG CAG TGG GGG ACA GTC TGT	917
Glu Gly Arg Leu Glu Val Tyr Tyr Lys Gly Gln Trp Gly Thr Val Cys	
265 270 275	
GAT GAT GGC TGG ACT GAG ATG AAC ACA TAC GTG GCT TGT CGA CTG CTG	965
Asp Asp Gly Trp Thr Glu Met Asn Thr Tyr Val Ala Cys Arg Leu Leu	
280 285 290	
GGA TTT AAA TAC GGC AAA CAG TCC TCT GTG AAC CAT TTT GAT GGC AGC	1013
Gly Phe Lys Tyr Gly Lys Gln Ser Ser Val Asn His Phe Asp Gly Ser	
295 300 305	
AAC AGG CCC ATA TGG CTG GAT GAC GTC AGC TGC TCA GGA AAA GAA GTC	1061
Asn Arg Pro Ile Trp Leu Asp Asp Val Ser Cys Ser Gly Lys Glu Val	
310 315 320 325	
AGC TTC ATT CAG TGT TCC AGG AGA CAG TGG GGA AGG CAT GAC TGC AGC	1109
Ser Phe Ile Gln Cys Ser Arg Arg Gln Trp Gly Arg His Asp Cys Ser	
330 335 340	
CAT AGA GAA GAT GTG GGC CTC ACC TGC TAT CCT GAC AGC GAT GGA CAT	1157
His Arg Glu Asp Val Gly Leu Thr Cys Tyr Pro Asp Ser Asp Gly His	
345 350 355	
AGG CTT TCT CCA GGT TTT CCC ATC AGA CTA GTG GAT GGA GAG AAT AAG	1205
Arg Leu Ser Pro Gly Phe Pro Ile Arg Leu Val Asp Gly Glu Asn Lys	
360 365 370	
AAG GAA GGA CGA GTG GAG GTT TTT GTC AAT GGC CAA TGG GGA ACA ATC	1253
Lys Glu Gly Arg Val Glu Val Phe Val Asn Gly Gln Trp Gly Thr Ile	
375 380 385	
TGC GAT GAC GGA TGG ACC GAT AAG CAT GCA GCT GTG ATC TGC CGG CAA	1301
Cys Asp Asp Gly Trp Thr Asp Lys His Ala Ala Val Ile Cys Arg Gln	
390 395 400 405	
CTT GGC TAT AAG GGT CCT GCC AGA GCA AGG ACT ATG GCT TAT TTT GGG	1349
Leu Gly Tyr Lys Gly Pro Ala Arg Ala Arg Thr Met Ala Tyr Phe Gly	
410 415 420	
GAA GGA AAA GGC CCC ATC CAC ATG GAT AAT GTG AAG TGC ACA GGA AAT	1397
Glu Gly Lys Gly Pro Ile His Met Asp Asn Val Lys Cys Thr Gly Asn	
425 430 435	

GAG AAG GCC CTG GCT GAC TGT GTC AAA CAA GAC ATT GGA AGG CAC AAC Glu Lys Ala Leu Ala Asp Cys Val Lys Gln Asp Ile Gly Arg His Asn 440 445 450	1445
TGC CGC CAC AGT GAG GAT GCA GGA GTC ATC TGT GAC TAT TTA GAG AAG Cys Arg His Ser Glu Asp Ala Gly Val Ile Cys Asp Tyr Leu Glu Lys 455 460 465	1493
AAA GCA TCA AGT AGT GGT AAT AAA GAG ATG CTC TCA TCT GGA TGT GGA Lys Ala Ser Ser Ser Gly Asn Lys Glu Met Leu Ser Ser Gly Cys Gly 470 475 480 485	1541
CTG AGG TTA CTG CAC CGT CGG CAG AAA CGG ATC ATT GGT GGG AAC AAT Leu Arg Leu Leu His Arg Arg Gln Lys Arg Ile Ile Gly Gly Asn Asn 490 495 500	1589
TCT TTA AGG GGT GCC TGG CCT TGG CAG GCT TCC CTC AGG CTG AGG TCG Ser Leu Arg Gly Ala Trp Pro Trp Gln Ala Ser Leu Arg Leu Arg Ser 505 510 515	1637
GCC CAT GGA GAC GGC AGG CTG CTT TGT GGA GCT ACC CTT CTG AGT AGC Ala His Gly Asp Gly Arg Leu Leu Cys Gly Ala Thr Leu Leu Ser Ser 520 525 530	1685
TGC TGG GTC CTG ACA GCT GCA CAC TGC TTC AAA AGG TAC GGA AAC AAC Cys Trp Val Leu Thr Ala Ala His Cys Phe Lys Arg Tyr Gly Asn Asn 535 540 545	1733
TCG AGG AGC TAT GCA GTT CGA GTT GGG GAT TAT CAT ACT CTG GTC CCA Ser Arg Ser Tyr Ala Val Arg Val Gly Asp Tyr His Thr Leu Val Pro 550 555 560 565	1781
GAG GAG TTT GAA CAA GAA ATA GGG GTT CAA CAG ATT GTG ATT CAC AGG Glu Glu Phe Glu Gln Glu Ile Gly Val Gln Gln Ile Val Ile His Arg 570 575 580	1829
AAC TAC AGG CCA GAC AGA AGC GAC TAT GAC ATT GCC CTG GTT AGA TTG Asn Tyr Arg Pro Asp Arg Ser Asp Tyr Asp Ile Ala Leu Val Arg Leu 585 590 595	1877
CAA GGA CCA GGG GAG CAA TGT GCC AGA CTA AGC ACC CAC GTT TTG CCA Gln Gly Pro Gly Glu Gln Cys Ala Arg Leu Ser Thr His Val Leu Pro 600 605 610	1925
GCC TGT TTA CCT CTA TGG AGA GAG AGG CCA CAG AAA ACA GCC TCC AAC Ala Cys Leu Pro Leu Trp Arg Glu Arg Pro Gln Lys Thr Ala Ser Asn 615 620 625	1973
TGT CAC ATA ACA GGA TGG GGA GAC ACA GGT CGT GCC TAC TCA AGA ACT Cys His Ile Thr Gly Trp Gly Asp Thr Gly Arg Ala Tyr Ser Arg Thr 630 635 640 645	2021
CTA CAA CAA GCT GCT GTG CCT CTG TTA CCC AAG AGG TTT TGT AAA GAG Leu Gln Gln Ala Ala Val Pro Leu Leu Pro Lys Arg Phe Cys Lys Glu 650 655 660	2069
AGG TAC AAG GGA CTA TTT ACT GGG AGA ATG CTC TGT GCT GGG AAC CTC Arg Tyr Lys Gly Leu Phe Thr Gly Arg Met Leu Cys Ala Gly Asn Leu 665 670 675	2117

- 32 -

CAA GAA GAC AAC CGT GTG GAC AGC TGC CAG GGA GAC AGT GGA GGA CCA	2165
Gln Glu Asp Asn Arg Val Asp Ser Cys Gln Gly Asp Ser Gly Gly Pro	
680 685 690	
CTC ATG TGT GAA AAG CCT GAT GAG TCC TGG GTT GTG TAT GGG GTG ACT	2213
Leu Met Cys Glu Lys Pro Asp Glu Ser Trp Val Val Tyr Gly Val Thr	
695 700 705	
TCC TGG GGG TAT GGA TGT GGA GTC AAA GAC ACT CCT GGA GTT TAT ACC	2261
Ser Trp Gly Tyr Gly Cys Gly Val Lys Asp Thr Pro Gly Val Tyr Thr	
710 715 720 725	
AGA GTC CCC GCT TTT GTA CCT TGG ATA AAA AGT GTC ACC AGT CTG	2306
Arg Val Pro Ala Phe Val Pro Trp Ile Lys Ser Val Thr Ser Leu	
730 735 740	
TAACTTATGG AAAGCTCAAG AAATAGTAAA ACAGTAACTA TTCAGTCTTC AAAAAAAAAA	2366
AAAAAAAAAA	2376

Patent claims**1. Neurotrypsins of the formulas I and II**

- 5 I: neurotrypsin of the human
 II: neurotrypsin of the mouse

 comprising the separate, coding and coded sequences of these compounds of the formulas I or II, comprising the separate partial sequences of the coding and coded sequences of these compounds of the formulas I or II, as for example the coding and coded sequences of the catalytic domains of the compounds of the formulas I or II, comprising the coding or coded sequences or partial sequences of the corresponding splice variants of the compounds of the formulas I or II, comprising the coding or coded sequences or partial sequences of the corresponding alleles of the compounds of the formulas I or II, comprising all sequence variants of the coding or coded sequences, or parts thereof, of the compounds of the formulas I or II, whose biological activity is equal or similar to that of the compounds of the formulas I or II, for example sequence variants of the compounds of the formulas I or II, which differ in the not conserved amino acid sequence positions of the sequence of the formulas I or II, comprising the sequences hybridizing to the coding sequences, or parts thereof, under stringent conditions, comprising the translation products of the sequences hybridizing to the coding sequences of the compounds of the formulas I or II, or to parts thereof, under stringent conditions, comprising the nucleotide sequences coding the proteins coded by the compounds of the formulas I or II, or parts thereof, but, as a result of the use of different alternative codons, are degenerated with regard to the nucleotide sequences defined by the compounds of the formulas I or II.

2. Pharmaceutical composition, characterized in that it contains as at least one active compound either the coded sequence or the coding sequence of the compound of the formula I or of the formula II, or the separate partial sequences of the coded and coding sequences of these compounds of the formulas I or II, as for example the coding or coded sequences of the catalytic domains, comprising the coding or coded sequences or partial sequences of the corresponding splice variants of the compounds of the formulas I or II, comprising the coding or coded sequences or partial sequences of the corresponding alleles of the compounds of the formulas I or II, comprising all

sequence variants of the coding or coded sequences, or parts thereof, of the compounds of formulas I or II, whose biological activity is equal or similar to that of the compounds of the formulas I or II, for example sequence variants of the compounds of the formulas I or II, which differ in the not conserved amino acid sequence positions of the sequence of the formulas I or II, comprising the sequences hybridizing to the coding sequences, or parts thereof, under stringent conditions, comprising the translation products of the sequences hybridizing to the coding sequences of the compounds of the formulas I or II, or to parts thereof, under stringent conditions, comprising the nucleotide sequences coding the proteins coded by the compounds of formulas I or II, or parts thereof, but, as a result of the use of different alternative codons, are degenerated with regard to the nucleotide sequences defined by the compounds of the formulas I or II.

3. Pharmaceutical composition, characterized in that it contains as at least one active compound a substance which changes the function of the coded sequence of the compounds of formulas I or II, for example, in that it reduces or increases the catalytic activity of the coded protein, or a part thereof, or in that it shortens or prolongs the time of presence of the coded protein at its place of action in the body.

4. Pharmaceutical composition, characterized in that it contains as at least one active compound a substance which changes the expression of the coding or coded sequences of the compounds of formulas I or II, for example in that it enhances or inhibits the transcription of the mRNA or in that it enhances or inhibits the translation of the coded sequences of the compounds of formulas I or II.

5. Pharmaceutical composition according to claim 2, 3, or 4, characterized in that it prevents or reduces the growth, the expansion, the infiltration and the metastasis of primary and metastatic tumors, as for example brain tumors or tumors of the retina.

6. Pharmaceutical composition according to claim 2, 3, or 4, characterized in that it contributes to the minimization of the tissue destruction in stroke, including brain infarction and ischemia, intracerebral hemorrhage, and subarachnoid hemorrhage, as for example by exerting a protecting effect on the cells of the so-called penumbra zone surrounding the necrotic tissue.

7. Pharmaceutical composition according to claim 2, 3, or 4, characterized in that it contributes to the minimization of the tissue destruction in traumatic brain injury, as for example by exerting a protective effect on the cells of the so-called zone surrounding the necrotic tissue.

5

8. Pharmaceutical composition according to claim 2, 3, or 4, characterized in that it prevents, ameliorates or cures the negative effects caused by neurodegenerative diseases.

10

9. Pharmaceutical composition according to claim 2, 3, or 4, characterized in that it prevents, ameliorates or cures the negative effects caused by neuroinflammatory diseases, as for example multiple sclerosis.

15

10. Pharmaceutical composition according to claim 2, 3, or 4, characterized in that it reduces or prevents negative effects on brain tissue caused by epileptic seizures.

20

11. Pharmaceutical composition according to claim 2, 3, or 4, characterized in that it contributes to the rescue of endangered neurons, as for example neurons endangered by hypoxia and ischemia, axotomy, nerve transection, deafferentiation, excitotoxicity, neuroinflammatory diseases and processes, epileptic seizures, and cancerous neoformations.

25

12. Pharmaceutical composition according to claim 2, 3, or 4, characterized in that it contributes to axonal regeneration and/or restoration of synaptic integrity and functions.

30

13. Pharmaceutical composition according to claim 2, 3, or 4, characterized in that it prevents, ameliorates, or cures retinal disorders, as for example retinal degeneration and retinal neoangiogenesis.

14. Pharmaceutical composition according to claim 2, 3, or 4, characterized in that it prevents cell death, comprising apoptosis and other forms of cell death, in the nervous system.

15. Pharmaceutical composition according to claim 14, characterized in that the cell death is an cell death in connection with damages of the nervous tissue, for example infarct of the brain and ischemic stroke, or hemorrhage of the brain, or trauma of the brain.

5

16. Pharmaceutical composition according to claim 14, characterized in that the cell death is an cell death in connection with damages of the nervous tissue, which occur due to lack of oxygen or glucose or due to intoxication.

10

17. Pharmaceutical composition according to claim 14, characterized in that the cell death is an cell death in connection with epileptic seizures.

15

18. Pharmaceutical composition according to claim 14, characterized in that the cell death is an cell death in connection with neurodegenerative diseases and inherited genetic deficiencies of the nervous system.

20

19. Pharmaceutical composition according to claim 14, characterized in that the cell death is an cell death in connection with axotomy or nerve transection, or deafferentiation.

25

20. Pharmaceutical composition according to claim 2, 3, or 4, characterized in that it influences the regeneration of injured, damaged, underdeveloped, or maldeveloped brain tissue and/or nervous tissue.

21. Pharmaceutical composition according to claim 2, 3, or 4, characterized in that it enhances the reorganization of the brain or nervous areas that have remained intact after brain and/or nerve injuries or after the destruction or damage of brain areas.

30

22. Pharmaceutical composition according to claim 2, 3, or 4, characterized in that it prevents, ameliorates, or cures pathological pain syndromes.

35

23. Pharmaceutical composition according to claim 2, 3, or 4, characterized in that it contributes to the improvement of the brain performance in healthy persons, as well as in persons with reduced brain performance.

24. Pharmaceutical composition according to claim 2, 3, or 4, characterized in that it ameliorates the learning and memory functions in healthy persons, as well as in persons with reduced learning and memory functions.

5 25. Pharmaceutical composition according to claim 2, 3, or 4, characterized in that it ameliorates or cures disorders in the field of disorders of the psychic wellness, or the psychosomatic state of health, as for example nervousity or „inner unrest“.

10 26. Pharmaceutical composition according to claim 2, 3, or 4, characterized in that it ameliorates or cures disorders in the field of the emotional functions, as for example states of anxiety.

 27. Pharmaceutical composition according to claim 2, 3, or 4, characterized in that it ameliorates or cures psychiatric disorders.

15

 28. Pharmaceutical composition according to claim 27, characterized in that the psychiatric disorder is a disorder in the field of schizophrenia and schizophrenia-like disorders, comprising chronic schizophrenia, chronic schizo-affective disorders, unspecific disorders, comprising acute and chronic schizophrenia of various
20 symptomatologies, as for example severe, non-remitting „Kraepelinic“ schizophrenia, or as for example the DSM-III-R-prototype of the schizophrenia-like disorders, comprising episodic schizophrenic disorders, comprising delusional schizophrenia-like disorders, comprising schizophrenia-like personality disorders, as for example schizophrenia-like personality disorders with mild symptomatics, comprising schizotypic personality
25 disorders, comprising the latent forms of schizophrenic or schizophrenia-like disorders, comprising non-organic psychotic disorders.

 29. Pharmaceutical composition according to claim 27, characterized in that the psychiatric disorder is a disorder in the field of the endogenic depressions or in the field
30 of manic or manic-depressive disorders.

 30. Pharmaceutical composition according to claim 2, 3, or 4, characterized in that it ameliorates or cures disorders of the brain function due deficiency, malfunction, or overfunction of at least one protease.

35

31. Pharmaceutical composition according to claim 30, characterized in that the protease is tissue-type plasminogen activator, abbreviated as tPA, urokinase-type plasminogen activator, abbreviated as uPA, or plasmin.

5 32. Pharmaceutical composition according to claim 2, 3, or 4, characterized in that it ameliorates or cures disorders of the function of the lungs due to deficiency, malfunction, or overfunction of at least one protease.

10 33. Pharmaceutical composition according to claim 32, characterized in that the disorder of the function of the lungs is chronic bronchitis or emphysema of the lungs.

34. Use for the production of recombinant proteins of the coding nucleotide sequences of the compounds of the formulas I or II, comprising the separate partial sequences of the coding sequences of the compounds of the formulas I or II, as for
15 example the coding sequences of the catalytic domains of the compounds of the formulas I or II, comprising the coding nucleotide sequences or partial sequences of the corresponding splice variants of the compounds of the formulas I or II, comprising the coding sequences or partial sequences thereof of the corresponding alleles of the compounds of the formulas I or II, comprising all sequence variants of the coding
20 sequences, or parts thereof, of the compounds of formulas I or II, whose translation products have a biological activity equal or similar to that of the translation products of the compounds of the formulas I or II, for example sequence variants of the compounds of the formulas I or II, which differ in the not conserved amino acid sequence positions of the sequence of the formulas I or II, comprising the sequences hybridizing to the coding
25 sequences of the compounds of the formulas I or II, or parts thereof, under stringent conditions, comprising the nucleotide sequences coding the proteins coded by the compounds of the formulas I or II, or parts thereof, but, as a result of the use of different alternative codons, are degenerated with regard to the nucleotide sequences defined by the compounds of the formulas I or II.

30

35. Use as targets for the development of pharmaceutical drugs, for example for the inhibition or the enhancement of the catalytic activity of the coded proteins of the formulas I or II, of proteins with the coded amino acid sequences of the compounds of the formulas I or II, comprising the proteins with the separate partial sequences of the
35 coded amino acid sequences of the compounds of the formulas I or II, as for example

the separate catalytic domains of the compounds of the formulas I or II, comprising the proteins with the coded sequences or partial sequences of the corresponding splice variants of the compounds of the formulas I or II, comprising the proteins with the coded amino acid sequences or partial sequences thereof of the corresponding alleles of the compounds of the formulas I or II, comprising all sequence variants of the coded sequences, or parts thereof, of the compounds of formulas I or II, whose biological activity is equal or similar to the coded sequences of the compounds of the formulas I or II, for example sequence variants of the compounds of the formulas I or II, which differ in the not conserved amino acid sequence positions of the sequences of the formulas I or II, comprising the proteins with the coded amino acid sequences, or partial sequences thereof, of the nucleotide sequences hybridizing to the coding sequences of the compounds of the formulas I or II, or parts thereof, under stringent conditions.

36. Use as targets for the development of pharmaceutical drugs, for example for the enhancement or the inhibition of the catalytic activity of the coded proteins of the formulas I or II, of the species-homologous proteins, or parts thereof, of the compounds of the formulas I or II, as for example the species-homologous proteins of the rat, the rabbit, the cow, the sheep, the pig, the primates, the birds, the zebra fish, the fruit fly (*Drosophila melanogaster*), etc., comprising the partial sequences thereof, as for example the separate catalytic domains, comprising the splice variants of the species-homologous proteins, comprising the alleles of the species-homologous proteins, comprising the translation products of the sequences hybridizing under stringent conditions to the corresponding species-homologous compounds of the formulas I or II, or their splice variants, or their alleles, of the coding sequences or partial sequences of the compounds of formulas I or II.

37. Use for the spatial structure determination, for example the spatial structure determination by means of crystallography or nuclear resonance spectroscopy, of the proteins with the coded amino acid sequences of the compounds of the formulas I or II, comprising the proteins with the separate partial sequences of the coded amino acid sequences of the compounds of the formulas I or II, as for example the separate catalytic domains, comprising the proteins with the coded sequences or partial sequences of the corresponding splice variants of the compounds of the formulas I or II, comprising the proteins with the coded amino acid sequences, or partial sequences thereof, of the corresponding alleles of the compounds of the formulas I or II, comprising

all sequence variants of the coded sequences, or parts thereof, of the compounds of the formulas I or II, whose biological activity is equal or similar to that of the coded sequences of the compounds of the formulas I or II, for example sequence variants of the compounds of the formulas I or II, which differ in the not conserved amino acid sequence positions of the sequences of the formulas I or II, comprising the translation products with the sequences hybridizing to the coding sequences of the compounds of the formulas I or II, or parts thereof, under stringent conditions, comprising the species-homologous proteins of the compounds of the formulas I or II, for example the species-homologous proteins of the rat, the rabbit, the cow, the sheep, the pig, the primates, the birds, the zebra fish, the fruit fly (*Drosophila melanogaster*), etc., comprising the partial sequences thereof, as for example the separate catalytic domains.

38. Use for the prediction of the protein structure by means of computerized protein structure prediction methods, of the coded amino acid sequences of the compounds of the formulas I or II, comprising the separate partial sequences of the coded amino acid sequences of the compounds of the formulas I or II, as for example the coded amino acid sequences of the separate catalytic domains of the compounds of the formulas I or II, comprising the coded sequences or partial sequences of the corresponding splice variants of the compounds of the formulas I or II, comprising the coded amino acid sequences, or parts thereof, of the corresponding alleles of the compounds of the formulas I or II, comprising all sequence variants of the coded sequences, or parts thereof, of the compounds of the formulas I or II, whose biological activity is equal or similar to that of the coded sequences of the compounds of the formulas I or II, for example sequence variants of the compounds of the formulas I or II, which differ in the not conserved amino acid sequence positions of the sequences of the formulas I or II, comprising the amino acid sequences of the translation products of the sequences hybridizing to the coding sequences of the compounds of the formulas I or II, or parts thereof, under stringent conditions, comprising sequences of the species-homologous compounds of the compounds of the formulas I or II, for example the sequences of the species-homologous compounds of the rat, the rabbit, the cow, the sheep, the pig, the primates, the birds, the zebra fish, the fruit fly (*Drosophila melanogaster*), etc., comprising the partial sequences of the species-homologous compounds, as for example the sequences of the catalytic domains of the species-homologous compounds.

35

39. Use as targets for the development of pharmaceutical drugs, for example for the inhibition or the enhancement of the catalytic activity of the coded proteins of the compounds of the formulas I or II, of the spatial structure of the coded amino acid sequences of the compounds of the formulas I or II, comprising the spatial structures of the separate partial sequences of the compounds of the formulas I or II, as for example the spatial structure of the catalytic domains, comprising the spatial structure of the coded sequences or partial sequences of the corresponding splice variants of the compounds of the formulas I or II, comprising the spatial structure of the coded sequences or partial sequences of the corresponding alleles of the compounds of the formulas I or II, comprising the spatial structure of all sequence variants of the coded sequences, or parts thereof, of the compounds of formulas I or II, whose biological activity is equal or similar to the coded sequences of the compounds of the formulas I or II, for example sequence variants of the compounds of the formulas I or II, which differ in the not conserved amino acid sequence positions of the sequences of the formulas I or II, comprising the spatial structures of the translation products of the sequences hybridizing to the coding sequences of the compounds of the formulas I or II, or parts thereof, under stringent conditions, comprising the spatial structures of the species-homologous compounds of the compounds of the formulas I or II, as for example the spatial structures of the species homologous compounds, or parts thereof, of the rat, the rabbit, the cow, the sheep, the pig, the primates, the birds, the zebra fish, the fruit fly (*Drosophila melanogaster*), etc..

40. Use in gene therapeutical applications in humans and in animals, as for example as parts of gene therapy vectors or as for example as parts of artificial chromosomes, of the coding nucleotide sequences of the compounds of the formulas I or II, comprising the separate partial sequences of the coding sequences of these compounds of the formulas I or II, as for example the coding sequences of the catalytic domains of the compounds of the formulas I or II, comprising the coding sequences or partial sequences of the corresponding splice variants of the compounds of the formulas I or II, comprising the coding sequences or partial sequences of the corresponding alleles of the compounds of the formulas I or II, comprising all sequence variants of the coding sequences, or parts thereof, of the compounds of the formulas I or II, whose translation products exhibit a biological activity which is equal or similar to that of the translation products of the compounds of the formulas I or II, for example sequence variants of the compounds of the formulas I or II, which differ in the not conserved amino

acid sequence positions of the sequences of the compounds of the formulas I or II, comprising the sequences hybridizing to the coding sequences, or parts thereof, under stringent conditions, comprising the nucleotide sequences coding the proteins coded by the compounds of the formulas I or II, or parts thereof, but as a result of the use of different alternative codons, are degenerated with regard to the nucleotide sequences defined by the compounds of the formulas I or II.

41. Use for so-called cell engineering applications for the production of gene technologically mutated cells, which produce the coded sequences, or parts thereof, of the compounds of the formulas I or II, for example for cell-therapeutical applications as a pharmaceutical composition according to claim 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, or 33, of the coding nucleotide sequences of the compounds of the formulas I or II, comprising the separate partial sequences of the coding sequences of these compounds of the formulas I or II, as for example the coding sequences of the catalytic domains of the compounds of the formulas I or II, comprising the coding sequences or partial sequences of the corresponding splice variants of the compounds of the formulas I or II, comprising the coding sequences or partial sequences of the corresponding alleles of the compounds of the formulas I or II, comprising all sequence variants of the coding sequences, or parts thereof, of the compounds of the formulas I or II, whose translation products exhibit a biological activity which is equal or similar to that of the translation products of the compounds of the formulas I or II, for example sequence variants of the compounds of the formulas I or II, which differ in the not conserved amino acid sequence positions of the sequence of the compounds of the formulas I or II, comprising the sequences hybridizing to the coding sequences, or parts thereof, under stringent conditions, comprising the nucleotide sequences coding the proteins coded by the compounds of formulas I or II, or parts thereof, but as a result of the use of different alternative codons, are degenerated with regard to the nucleotide sequences defined by the compounds of the formulas I or II.

42. Use as antigens for the production of antibodies, as for example antibodies that inhibit or promote the protease function or antibodies that can be used for immunohistochemical studies, of the coded amino acid sequences of the compounds of the formulas I or II, comprising the separate partial sequences of the coded amino acid sequences of the compounds of the formulas I or II, as for example the coded amino

acid sequence of the catalytic domain or one or more of the other domains or segments, comprising the coded sequences or partial sequences of the corresponding splice variants of the compounds of the formulas I or II, comprising the coded sequences or partial sequences of the corresponding alleles of the compounds of the formulas I or II, comprising all sequence variants of the coded sequences, or parts thereof, of the compounds of the formulas I or II, whose biological activity is equal or similar to that of the coded sequences of the compounds of the formulas I or II, for example sequence variants of the compounds of the formulas I or II, which differ in the not conserved amino acid sequence positions of the sequence of the compounds of the formulas I or II, comprising the translation products or parts thereof, of the sequences hybridizing to the coding sequences of the compounds of the formulas I or II, or parts thereof, under stringent conditions, comprising the coded sequences of the species-homologous compounds of the compounds of the formulas I or II, as for example the coded sequences of the species-homologous compounds of the rat, the rabbit, the cow, the sheep, the pig, the primates, the birds, the zebra fish, the fruit fly (*Drosophila melanogaster*), etc., comprising the separate partial sequences of the coded sequences of the species-homologous compounds of the compounds of the formulas I or II, as for example the coded amino acid sequence of the catalytic domain, or one or more of the other domains or segments.

20

43. Use for the production of transgenic animals, as for example transgenic mice, of the coding nucleotide sequences of the compounds of the formulas I or II, comprising the separate partial sequences of the coding sequences of these compounds of the formulas I or II, as for example the coding sequences of the catalytic domains of the compounds of the formulas I or II, comprising the coding sequences or partial sequences of the corresponding splice variants of the compounds of the formulas I or II, comprising the coding sequences, or partial sequences, of the corresponding alleles of the compounds of the formulas I or II, comprising all sequence variants of the coding sequences, or parts thereof, of the compounds of the formulas I or II, whose translation products exhibit a biological activity which is equal or similar to that of the translation products of the compounds of the formulas I or II, for example sequence variants of the compounds of the formulas I or II, which differ in the not conserved amino acid sequence positions of the sequences of the compounds of the formulas I or II, comprising the sequences hybridizing to the coding sequences, or parts thereof, under stringent conditions, comprising the nucleotide sequences coding the proteins coded by the

35

compounds of the formulas I or II, or parts thereof, but as a result of the use of different alternative codons, are degenerated with regard to the nucleotide sequences defined by the compounds of the formulas I or II.

5 44. Use for the inactivation or the mutation of the corresponding gene by means of gene targeting techniques, as for example the elimination of the gene in the mouse through homologous recombination, of the coding nucleotide sequences of the compounds of the formulas I or II, comprising the separate partial sequences of the coding sequences of these compounds of the formulas I or II, as for example the coding
10 sequences of the catalytic domains of the compounds of the formulas I or II, comprising the coding sequences, or partial sequences, of the corresponding splice variants of the compounds of the formulas I or II, comprising the coding sequences, or partial sequences, of the corresponding alleles of the compounds of the formulas I or II, comprising all sequence variants of the coding sequences, or parts thereof, of the
15 compounds of the formulas I or II, whose translation products exhibit a biological activity which is equal or similar to that of the translation products of the compounds of the formulas I or II, for example sequence variants of the compounds of the formulas I or II, which differ in the not conserved amino acid sequence positions of the sequence of the compounds of the formulas I or II, comprising the sequences hybridizing to the coding
20 sequences, or parts thereof, under stringent conditions, comprising the nucleotide sequences coding the proteins coded by the compounds of the formulas I or II, or parts thereof, but as a result of the use of different alternative codons, are degenerated with regard to the nucleotide sequences defined by the compounds of the formulas I or II.

25 45. Use for the diagnostics of disorders in the gene corresponding to the compound of the formula I, of the coding nucleotide sequences of the compounds of the formulas I or II, comprising the separate partial sequences of the coding sequences of these compounds of the formulas I or II, as for example the coding sequences of the catalytic domains of the compounds of the formulas I or II, comprising the coding
30 sequences or partial sequences of the corresponding splice variants of the compounds of the formulas I or II, comprising the coding sequences, or partial sequences, of the corresponding alleles of the compounds of the formulas I or II, comprising all sequence variants of the coding sequences, or parts thereof, of the compounds of the formulas I or II, whose translation products exhibit a biological activity which is equal or similar to that
35 of the translation products of the compounds of the formulas I or II, for example

sequence variants of the compounds of the formulas I or II, which differ in the not conserved amino acid sequence positions of the sequences of the compounds of the formulas I or II, comprising the sequences hybridizing to the coding sequences, or parts thereof, under stringent conditions, comprising the nucleotide sequences coding the proteins coded by the compounds of the formulas I or II, or parts thereof, but as a result of the use of different alternative codons, are degenerated with regard to the nucleotide sequences defined by the compounds of the formulas I or II.

46. Use as a starting sequence for gene technological modifications aimed at the production of pharmaceutical compositions or gene therapy vectors which exhibit changed properties as compared with the corresponding pharmaceutical compositions or gene therapy vectors containing the coding nucleotide sequence of the compounds of formulas I or II, for example changed proteolytic activity, changed proteolytic specificity, or changed pharmacokinetic characteristics, of the coding nucleotide sequences of the compounds of the formulas I or II, comprising the separate partial sequences of the coding sequences of these compounds of the formulas I or II, as for example the coding sequences of the catalytic domains of the compounds of the formulas I or II, comprising the coding sequences or partial sequences of the corresponding splice variants of the compounds of the formulas I or II, comprising the coding sequences, or partial sequences, of the corresponding alleles of the compounds of the formulas I or II, comprising all sequence variants of the coding sequences, or parts thereof, of the compounds of the formulas I or II, whose translation products exhibit a biological activity which is equal or similar to that of the translation products of the compounds of the formulas I or II, for example sequence variants of the compounds of the formulas I or II, which differ in the not conserved amino acid sequence positions of the sequences of the compounds of the formulas I or II, comprising the sequences hybridizing to the coding sequences, or parts thereof, under stringent conditions, comprising the nucleotide sequences coding the proteins coded by the compounds of the formulas I or II, or parts thereof, but as a result of the use of different alternative codons, are degenerated with regard to the nucleotide sequences defined by the compounds of the formulas I or II.

INTERNATIONAL SEARCH REPORT

national Application No

PCT/IB 98/00625

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/57 C12N9/64 A61K31/70 A61K38/48 C12Q1/37
 A61K48/00 C07K16/40 C12N15/00 A01K67/027 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A61K C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EMBL/GENBANK DATABASES Accession no AA166524, Sequence identification MMAA66524, 21 December 1996 MARRA M ET AL: "The WashU-HHMI Mouse EST Project" XP002075099 see the whole document	1, 34, 37, 38, 42
X	EMBL/GENBANK DATABASES Accession no AA373034, Sequence identification HSZZ78160, 18 April 1997 ADAMS M ET AL: "Initial assessment of human gene diversity and expression patterns based upon 83 million nucleotides of cDNA sequence" XP002075100 see the whole document	1, 34, 37, 38, 42



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
 "E" earlier document but published on or after the international filing date
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 "O" document referring to an oral disclosure, use, exhibition or other means
 "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

25 August 1998

Date of mailing of the international search report

04/09/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Van der Schaal, C

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 98/00625

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>EMBL/GENBANK DATABASES Accession no AA073513, Sequence identification MMA73513, 5 October 1996 MARRA M ET AL: "The WashU-HHMI Mouse EST Project" XP002075101 see the whole document</p>	1, 34, 37, 38, 42
X	<p>EMBL/GENBANK DATABASES Accession no AA063841, Sequence identification MMA63841, 25 September 1996 MARRA M ET AL: "The WashU-HHMI Mouse EST Project" XP002075102 see the whole document</p>	1, 34, 37, 38, 42
A	<p>YAMASHIRO K ET AL: "Molecular cloning of a novel trypsin-like serine protease (neurosin) preferentially expressed in brain." BIOCHIMICA ET BIOPHYSICA ACTA, (1997 JAN 3) 1350 (1) 11-4. JOURNAL CODE: AOW. ISSN: 0006-3002., XP002075096 Netherlands</p>	
P, X	<p>DATABASE MEDLINE US NATIONAL LIBRARY OF MEDICINE (NLM), BETHESDA, MD, US PROBA K ET AL: "Cloning and sequencing of the cDNA encoding human neurotrypsin." XP002075103 see abstract & BIOCHIMICA ET BIOPHYSICA ACTA, (1998 MAR 9) 1396 (2) 143-7. JOURNAL CODE: AOW. ISSN: 0006-3002., Netherlands</p>	1-4, 34-39, 42-44
P, X	<p>GSCHWEND T P ET AL: "Neurotrypsin, a novel multidomain serine protease expressed in the nervous system." MOLECULAR AND CELLULAR NEUROSCIENCES, (1997) 9 (3) 207-19. JOURNAL CODE: BID. ISSN: 1044-7431., XP002075097 United States see the whole document</p>	1-46

-/-

INTERNATIONAL SEARCH REPORT

I. International Application No

PCT/IB 98/00625

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>YAMAMURA Y ET AL: "Molecular cloning of a novel brain-specific serine protease with a kringle-like structure and three scavenger receptor cysteine-rich motifs." BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, (1997 OCT 20) 239 (2) 386-92. JOURNAL CODE: 9Y8. ISSN: 0006-291X., XP002075098 United States see the whole document</p> <p>-----</p>	<p>1-4, 34-39, 42-44</p>

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB 98/00625

Box I Observations where certain claims were found uns archabl (C ntinuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 40 42 44 and 45 are (partially) directed to a method of treatment of the human/animal body or to a diagnostic method practised on the human/animal body, the search has been carried out and based on the alleged effects of the compound.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.